

Observation and characterization of laser driven shock waves in tenuous plasma

H.AHMED¹, D.Doria¹, L. Romagnani², G.Sarri¹, E.Ianni¹, R. Prasad¹, M.Cerchez³,
A.L.Lindemann³, K.Quinn¹, O.Willi³, I. Kourakis¹, M. Borghesi¹

¹ *Centre for Plasma Physics, Queen's University Belfast, BT7 1NN, Belfast, Northern Ireland, UK*

² *Laboratoire LULI, Ecole Polytechnique, France*

³ *Institute for Laser and Plasma Physics, university of Dsseldorf, Germany*

The observation and characterization of collisionless shock waves generated in laser plasma interactions is presented. The shock waves are generated by long pulse ~ 1 ns, intense (10^{15}W/cm^2) laser irradiation of solid targets and are observed to propagate in a tenuous ($10^{13} - 10^{16} \text{cm}^{-3}$), non magnetized background plasma. These nonlinear entities are detected and characterized by employing a proton imaging technique, which allows the simultaneous detection of propagation velocity, width of the shock front and electrostatic field associated with the shock, with high spatial and temporal resolution. Ion Acoustic Solitons (IAS) were observed under certain conditions and as inferred from the reconstructed associated electric field profile. The variation of IAS velocity and width as a function of the ambient parameters was characterized. The data show an increase in velocity and decrease in width as the density of the background plasma (generated via photo ionization of controlled, low density gas in a gas cell) is increased. The experimental results are interpreted in the frame work of Korteweg-de Vries nonlinear wave description and in agreement with theoretical predications.

Collisionless shocks are one of the most interesting phenomena in plasma physics and play a fundamental role in many astrophysical phenomena such the generation of highly energetic particles and cosmic rays during supernova explosions [1, 2]. There are two possibilities of collisionless shock formation in unmagnetized plasma; electrostatic shocks (ES shocks) [3, 4] and weibel mediated shocks [5]; both types have not been explored completely in laboratory yet in astrophysical context. As a result of progress in laser technology, now it is possible to replicate the astrophysical conditions in laboratory and the study of these phenomena has received great importance in last decade [2]. The relation of collisionless shock waves with other non

linear entities (like Ion acoustic Solitons (IAS) [4], Ion acoustic waves [6], Phase space electron-hole [7], etc) also plays a vital role for many astrophysical phenomena [1]. Nowadays, ES shock can be created in Laboratory by interaction of high energy pulse with solid. The expansion of laser ablated plasma into tenuous plasma (created via photo-ionization, mainly by radiation from laser plasma [8]) generates ES shocks and had been observed by employing optical probes, which could not resolve the shock front and distinguish different typologies of shocks [3]. Recently, the ES shock has been observed by employing proton projection imaging (PPI) techniques, which resolve the shock front with simultaneous measurement of propagation velocity, associated electric field with high temporal and spatial resolution [4].

In this paper, we present the experimental observation of propagation of laser driven electrostatic shocks in tenuous plasma. These shocks are generated by the expansion of warm plasma into rarefied ionized background [8] and characterized by laser accelerated protons as a charged particle probe [9].

The experiment was carried out using the VULCAN laser system at RAL. The schematic of the experimental setup is shown in figure 1(a). The CPA pulse of




Fig. 1: (a) The schematic of experimental setup, with time of flight arrangement of proton probing (b) Proton Image of the interaction of nanosecond pulse with $50\mu\text{m}$ Au stripe, showing shock waves in laser ablating plasma, the arrow shows the direction of the laser (c) zoom of observed structure.

pulse duration $\sim 1\text{ps}$ focused by an off axis ($f/3$) parabola to an intensity $\sim 5 \times 10^{19}\text{W}/\text{cm}^2$, onto $20\mu\text{m}$ Au target (proton target), is used to produce proton probe beam. The long pulse of duration $\sim 1\text{ns}$ was focused onto a Au stripe target (inferred as shock target) of thickness $50\mu\text{m}$ and width $70\mu\text{m}$ at an angle of 45° with an intensity $\sim 5 \times 10^{15}\text{W}/\text{cm}^2$. The setup was enclosed in a gas cell filled by nitrogen at pressure between $10^{-1}\text{mbar} - 10^{-4}\text{mbar}$. The proton imaging technique was the main diagnostic employed to detect and characterize the shock waves propagating in low density plasma. Interferometry was employed as well to characterize the laser ablated

plasma. The distance between the proton target and the shock target was 4 mm and the distance between the shock target and the proton detector was $L = 3.6\text{cm}$, giving a projection magnification $M = (l + L)/l \simeq 10$. The proton beam, after having probed the plasma, was recorded on a stack of several layers of dosimetrically calibrated radio chromic Films (RCF). The multilayer arrangement of the RCF detector and the broad spectral content of the proton beam provided temporal multi-frame capabilities for the proton probing within a single laser shot. In the time of flight arrangement, the probing time can be calculated by $\tau = \tau_o + \left(\frac{l}{c} \times \left(\frac{2\epsilon_p}{m_p c^2}\right)^{-1/2}\right)$ where $v_p = \left(\frac{2\epsilon_p}{m_p c^2}\right)^{-1/2}c$ and $m_p c^2 \simeq 940\text{MeV}$ and τ_o is the optical delay between lasers, ϵ_p and m_p is the energy of protons and mass of proton respectively. Data exemplifying the features observed by PPI is shown in figure 1(b) and (c). As a rule of thumb the electric fields are directed from the regions of a lighter blue color compared to the background (zones of reduced probe proton flux) towards the regions of darker blue color (increased flux). A pronounced modulation in the probe proton density, revealing a strongly modulated field distribution is observed at $\sim 1\text{mm}$ from the target. This modulation is interpreted as a shock structure propagating in tenuous plasma. The shock has aver-




Fig. 2: (a) Proton density line out across the shock front (b) reconstructed electric field E , resembles with theoretically predicted electric field profile of the IAS [4].

age radius of curvature $\sim 700\mu\text{m}$ and thickness $\sim 60\mu\text{m}$ and expands with an approximately spherical symmetry from the target. The observed proton density modulation reveals an electric field which changes sign across the shock [4]. The proton density modulation can be related to deflecting transverse electric field by

$$\frac{\delta n_p}{n_p} \simeq \frac{-eL}{2\epsilon_p M} \nabla_{\perp 0} \cdot \int_b E_{\perp} dz \quad (1)$$

Where $\delta n_p = n_p - n_{pu}$, with n_p and n_{pu} are perturbed and unperturbed density, ϵ_p is the proton energy, M is magnification, L is the distance between interaction region and detector; and b is the region of non zero charge density crossed by protons given by $b = 2\sqrt{(R + \delta R/2)^2 - R^2}$ with radius of structure R and thickness δR . The proton density modulation profile is shown in figure 2, which qualitatively agrees with theoretical the expected pattern of IAS [4]. The numerical integration of equation (1) leads to the electric field profile at shock front




Fig. 3: (a) Variation of the shock velocity and (b) the shock width versus the tenuous plasma pressure.

is shown in figure 2(b), having a amplitude $3 \times 10^7 V/m$.

The propagation velocity of shock can be measured in single laser shot, because different shock front position in RCF layers corresponding to different probing times. The shock structure is moving with $8 \times 10^5 m/sec$ at 0.3mbar, but velocity varies for different pressure of background gas (i.e different plasma density when ionnized). The variation of velocity versus the background plasma density, is shown in figure3(a). It shows that the velocity of shock decreases with increase of the background plasma pressure. As the PPI technique resolves the shock front, the width of the shock structure can be measured. The width of the shock also varies with background plasma pressure as shown in figure 3(b).

The electrostatic shock launched by the expansion of dense plasma into tenuous plasma, has been observed and characterized by means of the proton projection imaging technique. The employment of the proton projection imaging allows the simul-

taneous measurement of the spatial profile, electric field distribution and propagation velocity of a shock with a high temporal (\sim ps) and spatial ($\sim \mu m$) resolution.

We acknowledge support of the CLF beam access funding from STFC. We also acknowledge the support from AWE plc and the ELI project.

References

- [1] R.Z.Sagdeev and C.F.Kennel., *Sci.Am.***264**, 106 (1991).
- [2] B. A. Remington *et al.*, *Science*, **284**, 1488 (1999); S.V. Bulanov *et al.*, *Eur. Phys. J. D.***555**, 483 (2009).
- [3] A. R. Bell *et al.*, *Phys. Rev. A.* **38**, 1363 (1998).
- [4] L. Romagnani *et al.*, *Phys. Rev. Lett.* **101**, 025004 (2008).
- [5] Y. Kuramitsu *et al.*, *Phys. Rev. Lett.* **106**, 175002 (2011).
- [6] H. Washimi and T. Taninti *et al.*, *Phys. Rev. Lett.* **17**, 996 (1966).
- [7] G. Sarri *et al.*, *Phys. Plasmas*, **17**, 010701 (2010).
- [8] S.O.Dean *et al.*, *Phys. Rev. Lett.* **27**, 487 (1971).
- [9] M. Borghesi *et al.*, *Plasma Phys. Control. Fusion*, **43**, A267 (2001); M. Borghesi *et al.*, *Plasma Phys. Control. Fusion*, **50**, 124040 (2008); L. Romagnani *et al.*, *laser Part.Beams*, **26**, 241 (2008).