

First escaping fast ion measurements in ITER-like geometry using an activation probe

G.Bonheure¹, M.Hult², A.Fenyvesi³, S.Äkäslompolo⁴, D.Carralero⁵, D.Degering⁶,
A.de-Vismes Ott⁷, M.Garcia-Munoz⁸, B.Gmeiner⁵, A.Herrmann⁵, M.Laubenstein⁹, G.Lutter⁵,
J. Mlynar¹⁰, H.W.Mueller⁵, V.Rohde⁵, W.Suttrop⁵, G.Tardini⁵ and the ASDEX Upgrade
Team⁵

¹ERM-KMS, B-1000 Brussels, Belgium, Partner in the Trilateral Euregio Cluster

²EC-JRC-IRMM, Retieseweg 111, B-2440 Geel, Belgium

³Nuclear Research Institute, Hungarian Academy of Sciences, H-4026 Debrecen, Bem tér
18/c, Hungary

⁴Association EURATOM-Tekes, Aalto University, Department of Applied Physics, P.O.Box
14100 FI-00076 AALTO, Finland

⁵Max-Planck-Institut für Plasmaphysik, EURATOM-Assoziation, D-85748 Garching,
Germany

⁶VKTA Rossendorf e.V., Postfach 510119, 01314 Dresden, Germany

⁷IRSN-LMRE, Bois des Rames, Bât. 501, 91400 Orsay, France

⁸Faculty of Physics, Seville University, Seville, Spain

⁹Laboratori Nazionali del Gran Sasso, S.S. 17/bis km 18+910, I-67010 Assergi (AQ), Italy

¹⁰Association Euratom-IPP.CR, CZ-182 21 Praha 8, Czech Republic

1. Introduction

More research is needed to develop suitable diagnostics for measuring alpha particle confinement in ITER[1] and techniques relevant for fusion reactor conditions need further development[2-5]. Based on nuclear reactions, the activation probe is a novel concept first tested in JET [6 -15]. It may offer a more robust solution for performing alpha particle measurements in ITER. This paper describes the first escaping fast ion measurements performed at ASDEX Upgrade (AUG) tokamak using an activation probe. A detailed analysis, outside the scope of this contribution, will be published in a journal paper.

2. AUG Activation probe

The probe consists of a graphite sample holder and cover with a slit (see in Fig.1). The sample holder has four slots spaced 90⁰ apart in which to place the samples. The samples arrangement, size and material, are presented in table 1. The probe is mounted on a manipulator slightly above the tokamak mid-plane. The geometry is similar to that foreseen for an ITER activation probe[11]. The distance between the first sample and the plasma separatrix is ~ 80 mm.








Figure 1. Pictures of the new AUG activation probe. a) Left: The cylindrical graphite cover with 4mm wide slit. b) Right: View of Inner sample holder and six samples placed in slot 1

We analyzed our samples using ultra-low level gamma-ray spectrometry (ULGS) at 1) IRMM (Institute for Reference Materials and Measurements) in the 225m deep underground laboratory HADES located at the Belgian nuclear centre SCK•CEN in Mol, Belgium [17] 2) Rossendorf, in the underground laboratory Felsenkeller located at a depth of 47m close to Dresden, Germany [18], 3) Laboratoire souterrain de Modane (LSM) [19] located circa 1700 m under Fréjus mountain, France and 4) Laboratori Nazionali del Gran Sasso (LNGS) in the 3800 water meter equivalent (mwe) low background counting facility located in the Gran Sasso nearby Assergi in Italy[20].

3. Results and discussion

Each sample (see fig.1 and tab.1) was exposed to one plasma except the last row that was kept for background reference. Two H-mode deuterium plasmas ($B_t = 2.5$ T, $I_p = 10^6$ A) with long flat-top phases and high DD fusion yield were performed. The measured 2.5 MeV neutron emission is $7.0 \cdot 10^{15}$ in #29226 and $1.1 \cdot 10^{16}$ in #29228. The fusion reactions are:



Two proton activation products – ${}^7\text{Be}$ and ${}^{89}\text{Zr}$ – were found. ${}^7\text{Be}$ is mostly produced by 3 MeV protons, see in reaction (2) above. ${}^{89}\text{Zr}$ was measured on the Y7 sample and is produced by the 14.7 MeV protons (1). We identified four neutron induced products – ${}^{90m}\text{Y}$, ${}^{47}\text{Ca}$, ${}^{88}\text{Y}$ and ${}^{48}\text{Sc}$. Due to its short half-life (3.19h), ${}^{90m}\text{Y}$ could only be measured on-site. Due to its low activity and relatively short half-life (4.54d), ${}^{47}\text{Ca}$ was measured on C4 sample only and through its daughter ${}^{47}\text{Sc}$ on other C samples. ${}^{88}\text{Y}$ is produced by 14 MeV neutrons (4) and ${}^{48}\text{Sc}$ is produced mostly by 2.5 MeV neutrons (2). The measured 3 MeV proton fluence

in units of 10^8 cm^{-2} are: B1: 5.1 ± 1.7 , B3: 8.3 ± 2.2 , B5: 5.1 ± 2.2 and B7: 5.5 ± 2.8 . A fit over six experimental $^{10}\text{B}(p,\alpha)$ cross section data and TENDL2011 is used for this work[28-33].

Table I. Left: Overview and location of samples. The color refers to which laboratory analysed the sample – see in text above - (light grey-LNGS, pink-Modane, black-Felsenkeller and light blue-IRMM). The sample size is $10 \times 10 \text{ mm}$ and thickness 2 mm . The materials: Boron carbide B: B_4C , Calcium fluoride C: CaF_2 and Yttrium orthovanadate Y: YVO_4 . **Table II.** Right: Detected activation products ordered by mass number

Sample row	1	2	3	4	RN ^a	PR ^b	Thr ^c	S ^d
Plasma number	29226	29227	29228	n/a				
CaF2	C1	C2	C3	C4	^7Be ($t_{1/2} = 53.22 \text{ d}$)	$^{10}\text{B}(p,\alpha)$	--	B1,B2,B3, B5,B6,B7
B4C	B1	B2	B3	B4	^{47}Ca ($t_{1/2} = 4.54 \text{ d}$)	$^{46}\text{Ca}(n,\gamma)$ $^{47}\text{Ca}(n, 2n)$	- 10.2	C1, C3,C4 C5,C6
CaF2	C5	C6	C7 no data	C8 no data	^{48}Sc ($t_{1/2} = 1.82 \text{ d}$)	$^{51}\text{V}(n, \alpha)$	2.1	Y1,Y2,Y3, Y5,Y6,Y7
B4C	B5	B6	B7	B8	^{88}Y ($t_{1/2} = 106.63 \text{ d}$)	$^{89}\text{Y}(n,2n)$	11	All Y
YVO4	Y1	Y2	Y3	Y4	^{89}Zr ($t_{1/2} = 3.27 \text{ d}$)	$^{89}\text{Y}(p,n)$	3.7	Y7
YVO4	Y5	Y6	Y7	Y8	$^{90\text{m}}\text{Y}$ ($t_{1/2} = 3.19 \text{ h}$)	$^{89}\text{Y}(n, \gamma)$	-	Y ^e

^a RN: Radionuclide found, ^b PR: Main production reaction, ^c Thr: Energy threshold in MeV for the reaction, ^d S: Sample, ^e Measured at IPP-Garching, integrated measurement of all samples

As shown in table II, ^7Be is detected on all B samples except B4 and B8. Although B4 and B8 were exposed to gamma and neutron radiation, they were not exposed to escaping fast ions. The absence of ^7Be on B4 and B8 further confirms that it is due to escaping protons. These results suggest a higher proton loss for plasma #29228. If the loss per source proton is constant for both plasmas due to near-identical magnetic fields, the increase may be due to the proton source increase which is $\sim 50\%$ higher in #29228 than #29226. The comparison with ASCOT code Monte Carlo calculations [21, 22] which is an important goal of these experiments will provide detailed answers. Please see in the accompanying paper [23]. The ^{89}Zr detection is a new interesting result. It must be produced from reactions (1) due to confined ^3He ions. A spectral line corresponding to $^{91\text{m}}\text{Nb}$, produced by the $^{89}\text{Y}(^3\text{He},n)$ reaction –no threshold- was found. However this reaction cross section is unknown and the observed line may also be a statistical fluctuation.

4. Conclusions Results from the first activation probe experiment performed at AUG are reported. For the first time, single plasmas are measured. Absolute measurements of

mid-plane alpha-like escaping 3 MeV protons were performed. These measurements are extremely valuable for the benchmark of ASCOT Monte Carlo calculations. We found some evidence for ³He induced activation but more research is needed to confirm this new result. The ³He burn-up products measurement is also a new result. Finally, our experimental technique could easily be optimized by using a modified optimized aperture and larger samples area. We also opened up the possibility at AUG to measure alpha particles from reactions (1) without the need for tritium.

Acknowledgements

The contributions of all team members of IRMM, Modane, Felsenkeller, Gran Sasso and ASDEX-Upgrade are gratefully acknowledged. This work, supported by the European Communities under the contract of Association between EURATOM and Belgian state, was carried out within the framework of the European Fusion Development Agreement. The views and opinions expressed herein do not necessarily reflect those of the European Commission.

References

- [1] Donne A.J.H et al 2007 Nucl.Fusion **47** S337-S384
- [2] Zweben S.J et al, 1999 Nucl.Fusion **40** 91
- [3] Boivin R. et al, 1992 Nucl.Fusion **33** 93 449
- [4] Bonheure G. et al 2006 PoS(FNDA2006)091. Proc. of Science. Trieste : SISSA (International School for Advanced Studies), 2005. ISSN 1824-8039.
[http://pos.sissa.it/archive/conferences/025/091/FNDA2006_091.pdf] [International Workshop on Fast Neutron Detectors and Applications. University of Cape Town, South Africa (ZA), 03.04.2006-06.04.2006]
- [5] Darrow D. et al, 1995 Rev.Sci.Instrum. **66** 476
- [6] Bonheure G. et al 2008 Fus.Sci.Technol. **53** 3 806
- [7] E. Wieslander et al, Nucl. Instr. Meth **A. 591** (2008) 383-393
- [8] Bonheure G. et al, Nucl.Fusion **52** (2012) 083004
- [9] Bonheure G. et al Rev.Sci.Instrum. **81**,10D331-1 (2010)
- [10] González de Orduña R et al, Nucl. Instr. Meth **A 632** (2011) 89–100
- [11] Bonheure G. et al, Fusion Eng. Des. **86** (2011) 1298
- [12] Gasparro J. et al 2006 Appl.Radiat.Isot. 64 (10-11) 1130
- [13] Bonheure G. et al Physica Scripta **75** (2007) 769-773
- [14] Bonheure G. et al, Rev. Sci. Instrum. 83, 10D318 (2012)
- [15] Etxebarria N. et al 1996 Proc. of the Second International k0-Users, Ljubljana, Slovenia. pp. 137-141
- [16] Bonheure G. et al, Rev.Sci.Instrum. **79**,1 (2008)
- [17] Hult M. et al 2000 Appl. Radiat. Isot. **53** (1-2) 225-229
- [18] M. Köhler et al 2009 Appl. Radiat. Isot. **67** 5, 736-740
- [19] R.Gurriaran et al, Nucl. Instr. Meth **A. 524** (2004) 264-272
- [20] Arpesella C. et al 1996 Appl.Radiat.Isotopes **47** 991
- [21] Kurki Suonio T. et al 2009 Nucl. Fusion **49** 095001
- [22] J.A Heikkinen et al 1995 Physics of Plasmas 2 3724-33
- [23] S. P1.501 et al, this conference
- [24] Batistoni P. et al 1987 Nucl.Fus **27** 6 1043
- [25] Breizman B.N and Sharapov S.E 2011 Plasma Phys. Control. Fusion 53 054001
- [26] Günter S. et al 2007 Nucl. Fusion **47** 920
- [27] G.Vayakis et al., Fusion Sci. Technol. 53(February 2008), 699–750
- [28] W.E.Burcham et al., Philosophical Magazine 40 (1949) 807
- [29] W.E.Burcham et al., Philosophical Magazine 41 (1950) 337.
- [30] G. G. Bach et al., Philosophical Magazine 46 (1955) 824.
- [31] S. P. Kalinin et al., Atomnaya Energiya 2 (1957) 171.
- [32] J. Szabó et al., Conf.on Nucl.Data for Sci.and Technol., Antwerp 1982, p.956.
- [33] A.J. Koning and D. Rochman, TENDL-2011: TALYS-based Evaluated Nuclear Data Library, Release date: December 29, 2011, Nuclear Research and Consultancy Group (NRG), Petten, The Netherlands, 2011