

## Impurity Transport Studies at Wendelstein 7-X by Means of X-ray Imaging Spectrometer Measurements

A. Langenberg<sup>1</sup>, N.A. Pablant<sup>2</sup>, A. Dinklage<sup>1</sup>, Th. Wegner<sup>1</sup>, P. Traverso<sup>3</sup>, O. Marchuk<sup>4</sup>, B. Geiger<sup>1</sup>, B. Buttenschön<sup>1</sup>, C. Brandt<sup>1</sup>, H. Thomsen<sup>1</sup>, M. Kubkowska<sup>5</sup>, A. Czarnecka<sup>5</sup>, S. Jabłoński<sup>5</sup>, U. Neuner<sup>1</sup>, N. Tamura<sup>6</sup>, J.L. Valesco<sup>7</sup>, J.A. Alonso<sup>7</sup>, A. Mollén<sup>1</sup>, D. Zhang<sup>1</sup>, R. Burhenn<sup>1</sup>, R.C. Wolf<sup>1</sup> and the W7-X team

<sup>1</sup> Max-Planck-Institut für Plasmaphysik, 17491 Greifswald, Germany

<sup>2</sup> Princeton Plasma Physics Laboratory, Princeton, NJ, USA

<sup>3</sup> Auburn University, Auburn, Alabama, USA

<sup>4</sup> Institut für Energie und Klimaforschung-IEK-4, Forschungszentrum Jülich, 52425 Jülich, Germany

<sup>5</sup> Institute of Plasma Physics and Laser Microfusion, Hery 23 St. 01-497 Warsaw, Poland

<sup>6</sup> National Institute for Fusion Science, 322-6, Oroshi-cho, Toki-City, Gifu 509-5292, Japan

<sup>7</sup> Laboratorio Nacional de Fusión, Asociación EURATOM-CIEMAT, Madrid, Spain

Due to non axis symmetric 3D magnetic fields, impurity transport in the hot plasma core in stellarators is fundamentally different to tokamaks. In view of reactor-like operation, understanding the impurity transport is a prerequisite for steady-state operation. These aspects motivate initial impurity transport studies in W7-X at previously - in optimized stellarators – unexplored, reactor-relevant collisionalities. New effects, like potential variations on flux-surfaces [1] or screening effects due to species dependent transport regimes [2] are examples for aspects which attracted recent interest. Spatio-temporal impurity emissivities were measured by the x-ray imaging spectrometers XICS [3] and HR-XIS [4], optimized to detect He-like impurity emission. These spectrometers provide measurements of the radial electric field [5] and also allow for a direct determination of diffusive and convective transport parameters  $D$  and  $v$  [6]. Therefore, impurity transport in various stellarator specific transport regimes can be studied. In this paper, a systematic parameter scan varying the electron cyclotron resonance (ECR) heating power and the electron density  $n_e$  has been carried out. Furthermore, the specific settings of the power deposition reveal a significant impact on impurity confinement time, possibly driven by changes in the radial electric field at very low collisionalities – uniquely addressable in large stellarators like W7-X. Experimental findings are compared to neoclassical theory [7] and modeled with the 1D transport analysis code STRAHL. The study aims to reveal the impact of aspects entering stellarator optimization (e.g. ripples, magnetic mirrors) on the impurity fluxes.

[1] J.M Garcia-Regana et al, Nucl. Fusion 57, 056004 (2017)

[2] P. Helander et al., Phys. Rev. Lett. 118, 155002 (2017)

[3] N.A. Pablant, M. Bitter, R. Burhenn *et al.* 41<sup>st</sup> EPS conference on Plasma Physics Berlin (2014)

[4] G. Bertschinger, W. Biel, H. Jaegers, and O. Marchuk, Rev. Sci. Instrum. **75** 3727 (2004)

[5] N.A. Pablant, A. Langenberg, S. Satake *et al.* 43<sup>rd</sup> EPS conference on Plasma Physics Leuven (2016)

[6] A. Langenberg, N.A. Pablant, O. Marchuk *et al.* Nucl. Fusion **57** 086013 (2017)

[7] A. Mollén, S. Newton, P. Helander *et al.* 21<sup>st</sup> Int. Stellarator Workshop, Kyoto, (2017) (invited)