

## Computational modeling of quasi-single helicity states in an RFP

A.S. Ware and C. Miele

*University of Montana, Missoula, Montana, USA*

This work explores the impact of boundary shaping on access into and out of quasi-single helicity states in reverse-field-pinch (RFP) plasmas. Experiments have shown that RFP plasmas can self-organize to a quasi-single helicity (QSH) equilibrium with a helical axis [1, 2]. These states have improved confinement and lower magnetic turbulence levels compared to a standard RFP plasma which has multiple helicities in the magnetic spectrum. The VMEC code can obtain similar equilibria with a helical axis and a symmetric boundary [3]. These equilibria all have circular, or nearly-circular cross-sections. In this work we analyze the VMEC input parameters that control access to QSH states and test the impact of 2D-shaping of the boundary on RFP equilibria. Particular attention is paid to the impact that shaping has on access to quasi-single helicity states. The effect of increasing elongation and triangularity are tested systematically. Increased elongation results in lower plasma current for the same safety factor profile and a larger radial excursion of the helical axis in a QSH state. Optimization of the boundary coefficients targeting an increased radial excursion of the helical axis is undertaken. Results will be presented.

### References

- [1] D.F. Escande, et al., Phys. Rev. Lett. 85, 1662 (2000)
- [2] L. Marrelli, L., et al. Phys. Plasmas 9, 2868 (2002)
- [3] J.D. Hanson, et al., Nucl. Fusion 53, 083016 (2013)