

High-quality gamma-rays driven by petawatt laser pulse in near-critical density plasmas

T. W. Huang, C. T. Zhou, L. B. Ju, H. Zhang, and S. C. Ruan

College of Engineering Physics, Shenzhen Technology University, Shenzhen 518118, China

The nonlinear synchrotron radiation of direct laser-accelerated electrons in near-critical density (NCD) plasmas recently has been proposed as a very efficient scheme to produce multi-MeV gamma-rays [1]. In this presentation, we demonstrate that by employing a plasma density channel, the divergence angle and transverse size of the gamma-rays can be much reduced [2]. In addition, we propose a highly efficient gamma photon emitter obtained by irradiating a not-so-intense laser pulse on a miniature plasma device consisting of a plasma lens and a plasma mirror [3]. In this novel scheme, brilliant gamma-rays with very high conversion efficiency (higher than 1%) and spectral intensity (higher than 10^9 photons/0.1%BW/s) can be achieved by employing currently available lasers with intensity of 10^{21} W/cm². The practical effects of different nanostructures in the plasma lens and the oblique laser incidence are also discussed in this scheme [4]. At last, a novel scheme by exploiting an intense Laguerre Gaussian laser pulse interacting with under-dense plasmas is also proposed to produce helical gamma-rays with very small divergence angle (less than 5°) and ultra-high brilliance ($\sim 10^{24}$ photons/s/mm²/mrad²/0.1%BW) at a laser intensity of 10^{22} W/cm² [5]. Such high-quality gamma-rays generated in these schemes would find applications in wide-ranging areas.

References

- [1] D. J. Stark et al., Phys. Rev. Letts. 116, 185003 (2016).
- [2] T. W. Huang et al., Appl. Phys. Letts. 110, 021102 (2017).
- [3] T. W. Huang et al., New J. Phys. 21, 013008 (2019).
- [4] T. W. Huang et al., Plasma Phys. Control. Fusion 60, 115006 (2018).
- [5] L. B. Ju, T. W. Huang et al., submitted (2019).